



THE MEDICAL ASSOCIATION

OF THAILAND

Association of Medical Professionals in the
Kingdom of Thailand

Volume 31
No. 2
FEBRUARY
2008
ISSN 0125-2208



Co-Editors in Charge of

- The Medical Association of Thailand
- The Royal Thai College of Obstetricians and Gynecologists
- The Royal College of Physicians of Thailand
- Faculty of Medicine, Prince of Songkla University
- Commission on Higher Education
- Bureau of Personnel Administration and Development
- Faculty of Medicine, Mahasarakham University

***Pueraria Mirifica*, Phytoestrogen-Induced Change in Synaptophysin Expression Via Estrogen Receptor in Rat Hippocampal Neuron**

Rattiya Chindewa MSc^{*,**}, Saitida Lapanantasin^{*,**},
Yupin Sanvarinda PhD^{*,***}, Sukumal Chongthammakun PhD^{*,**}

* Center for Neuroscience, Faculty of Science, Mahidol University, Bangkok

** Department of Anatomy, Faculty of Science, Mahidol University, Bangkok

*** Department of Pharmacology, Faculty of Science, Mahidol University, Bangkok

Objective: To examine *Pueraria mirifica* (Leguminosae) containing-phytoestrogen effect on synaptic density and involvement of estrogen receptor.

Material and Method: The level of synaptophysin, a presynaptic vesicle protein, was measured using Western blot analysis and immunocytochemistry in hippocampal primary cell cultures at 6 days in vitro.

Results: *P. mirifica* and 17 β -estradiol (0.1 μ M) treatment for 4 days, but not for 2 days, significantly increased synaptophysin immunoreactivity and level of synaptophysin. *P. mirifica* up to 60 μ g/ml resulted in a dose related increase in the level of synaptophysin immunoreactivity. The classical estrogen receptor antagonist, ICI 182 780, significantly blocked *P. mirifica*-induced increase in synaptophysin.

Conclusion: *P. mirifica*-containing phytoestrogen affects synaptic density by inducing synaptophysin expression via estrogen receptor.

Keywords: Phytoestrogens, *Pueraria mirifica*, Hippocampal neuron, 17 β -estradiol, Synaptophysin, Synaptic density

J Med Assoc Thai 2008; 91 (2): 208-14

Full text. e-Journal: <http://www.medassocthai.org/journal>

Hippocampal neurons, which carry both estrogen receptor alpha (ER α) and estrogen receptor beta (ER β), are a very useful model for investigating estrogenic effect on the central nervous system⁽¹⁾. Dissociated cultures of hippocampal neurons, which have functional synapses⁽²⁾ and relative sparseness of dendritic spines⁽³⁾, offer a viable and convenient means for studying factors that regulate the development and function of dendritic spines or synapse formation^(3,4). A number of studies have suggested that synaptophysin, a 38-kDa calcium-binding glycoprotein, participates in synaptic vesicle formation, trafficking, release and recycling of vesicles containing neurotransmitters^(5,6). An increase in synaptophysin immunoreactivity has most often been interpreted as an increase in

presynaptic terminals reflecting increase in synapse formation⁽⁷⁾.

Estrogens have been reported to have cognitive enhancing and neuroprotective effects. They have been known to increase spine formation, synaptogenesis and synaptic proteins in hippocampal CA1 pyramidal neurons that mediate learning and memory⁽⁸⁾. The effects of estrogen replacement on memory and synaptophysin expression in aged post-estropausal female rodents suggest that it might be related to memory and synaptic plasticity as shown by increasing in synaptophysin expression⁽⁹⁾.

Phytoestrogens, plant-derived estrogen-like substances, have been extensively studied for their potential beneficial effects against hormone-dependent and age-related disease⁽¹⁰⁻¹²⁾. However, limited data exists regarding the effects of phytoestrogens on the brain. *Pueraria mirifica* (*P. mirifica*), is a Thai indigenous herb that belongs to the Family Leguminosae

Correspondence to : Chongthammakun S, Center for Neuroscience and Department of Anatomy, Faculty of Science, Mahidol University, Bangkok 10400, Thailand. Phone: 0-2201-5041, E-mail: scsct@mahidol.ac.th

with a long history of domestic consumption as a rejuvenating herb to promote youthfulness in both women and men⁽¹³⁾. Active phytoestrogenic ingredients found in its roots are isoflavones (daidzin, daidzein, genistin, genistein, and puerarin), and miroestrol and its derivatives including β -sitosterol, stigmasterol and coumestrol^(14,15). *P. mirifica* has estrogenic effects on ovarian function and the menstrual cycle in rats⁽¹⁶⁾ and monkeys⁽¹⁷⁾.

In order to examine whether *P. mirifica*-containing phytoestrogens affect synapse formation, the authors examined the role of *P. mirifica* on synaptophysin expression in primary hippocampal neuron cultures. The involvement of estrogen receptors with *P. mirifica* effects was also investigated.

Material and Method

Animals

Nineteen-old pregnant Sprague-Dawley rats were obtained from the National Laboratory Animal Center, Thailand. Experiments were carried out in accordance with the European Communities Council Directive of 24 November 1986 (86/609/EEC).

Primary cultures of prenatal hippocampal neurons

Primary cultures of prenatal hippocampal neurons were prepared from 19-day old Sprague-Dawley rats. Dissection of hippocampi, which were removed from anesthetized embryos, was done in dissociation media (0.08 M Na₂SO₄, 0.03 M K₂SO₄, 6 mM MgCl₂, 0.25 mM CaCl₂, 1 mM HEPES pH 7.4, 0.02 M glucose, phenol red), containing 1mM kynurenate, 10 mM MgCl₂ and 0.35 M NaOH. Hippocampi were digested with 1 mg/ml papain (Sigma, St. Louis, MO). Mechanical trituration was performed through a 5 ml pipette in L17 GM solution (0.2% NaHCO₃, 6.5% fetal calf serum, 35 mM glucose, 0.5 mM L-glutamine and 100 unit/ml pen/strep, 0.1 mM putrescine, 0.02M B 27 (GIBCO, Carlsbad, CA), 1X ITS (Sigma, St. Louis, MO), pH 7.4). About 10⁵ suspended cells/ml were plated onto 12 mm glass coverslips precoated with 2 μ g/ml of poly-L-lysine (Sigma, St. Louis, MO) and 2 μ g/ml of laminin (GIBCO, Carlsbad, CA) in 35 mm dishes for immunocytochemistry or at about 10⁶ cells/ml in 60 mm dishes for immunoblot. Cells were cultured at 37°C in 5% (v/v) CO₂/95% air atmosphere. During the first day of culture, 2.4 μ M cytosine arabinoside (Sigma, St. Louis, MO) were added to the cultures to prevent glia proliferation. Medium was changed to fresh L17 GM supplemented with B-27 after plating for two and a half hours⁽¹⁸⁾.

Preparation of *P. mirifica*

Tuberous roots of *P. mirifica* were collected in Chiang Mai province, Thailand. Plant materials were washed, dried, sliced into pieces, and ground to powder. Two g of plant powder were subsequently extracted with 250 ml of ethyl acetate for 4 h in a reflux apparatus. The eluted solution was evaporated to dryness with a rotary evaporator. Thirty mg of the dry brownish herb powder were dissolved in 1 ml of absolute ethanol overnight. The solution was filtered to remove undissolved powder and passed through a 0.22 μ m Millipore filter. The sterile filtrate was kept at -20°C before use.

Cell dosing

Experimental solutions were prepared on the day of dosing by adding 17 β -estradiol (Sigma, St. Louis, MO) and *P. mirifica* in growth media to give a final concentrations of 0.1 μ M and 30 μ g/ml respectively on the second and fourth day of culture. Cultured cells (treated and untreated) were collected on the 6th day *in vitro* and were then processed for immunocytochemistry or immunoblot. For treatment with the estrogen antagonist, 1 μ M ICI 182 780 (Tocris, Bristol, England) was added prior to adding *P. mirifica*.

Immunocytochemistry

The 6-day *in vitro* treated and untreated hippocampal neurons on cover slips were washed with 0.01 M phosphate buffer saline (PBS) once, then fixed with 100% methanol at -20°C for 10-15 min. Autofluorescence of cells was quenched twice, 5 min each time, with 10 mM glycine in PBS and permeabilized with 0.5% NP40 in PBS for 5 min. After treating with blocking solution (10% normal goat serum (NGS), 1% bovine serum albumin (BSA), 0.5% Triton X-100 in PBS) for 1 h, cells were incubated for 1 h at room temperature with rabbit primary anti-human synaptophysin antibody (Zymed, South San Francisco) diluted 1:250 in primary antibody buffer (3% NGS, 1% BSA, 0.5% Triton X-100, 0.05% sodium azide in PBS). After washing to remove excess antibodies with PBS, cells were incubated at room temperature with secondary goat anti-rabbit IgG TRITC-conjugated antibody (Zymed, South San Francisco, CA) diluted 1:500 in secondary antibody buffer (3% NGS, 1% BSA, 0.5% Triton X-100 in PBS) for 1 h. Cells were washed three times with PBS, mounted with vectashield mounting media and examined under light and fluorescence microscopy with a 40x objective lens and using laser at excitation of 553 nm.

Western blot

Treated and untreated neurons were washed with PBS three times then lysed in 100 μ l of lysis buffer (50 mM Tris pH 7.4, 150 mM NaCl, 1% Triton X-100, 1% DOC, 0.1% SDS, 5 mM EDTA, 30 mM Na_2HPO_4 , 50 mM NaF, 0.5 mM NaVO_4 , 2 mM PMSF, 1% aprotinin and 1% leupeptin). per 60 mm dish. Insoluble materials were removed by high-speed centrifuge at 12,000xg at 4°C for 10 min. The supernatants were carefully placed into clean tubes, and protein concentrations were determined by the Bradford protein assay. Equal amounts of protein (10 μ g per lane) in the loading buffer were denatured at 95°C for 5 min and separated by 12.5% SDS-PAGE. Proteins in the gels were transferred to a nitrocellulose membrane, washed with TBST (1 M Tris pH 7.4, 3 M NaCl, 0.1% Tween 20) and incubated with 5% skim milk in TBST. Membranes were stored at 4°C

overnight and then were incubated with primary antisynaptophysin antibody (1:1000 in TBST). After washing three times with TBST, the membranes were incubated with a horseradish peroxidase-conjugated secondary antibody (1:5000 in TBST) (Zymed, South San Francisco, CA) and treated using a LumiGLO Chemiluminescent Substrate kit (KPL, Gaithersburg, MD).

Quantification

The number of radiant dots, shown in Fig. 1A, was used as a measure of the level of synaptophysin immunoreactivity. The number of dots was randomly counted per 50 μ m of neuronal process. Manual counting was performed by a person unaware of the treatment protocol. Five to ten processes were counted per image. At least 3 cover slips per group were processed, at least 3 images were taken for each cover




Fig. 1 Effect of *P. mirifica* on synaptophysin expression in rat hippocampal neurons
 (A) Immunocytochemical detection of synaptophysin in 6-day cultured hippocampal neuron treated with 0.1 μ M 17 β -estradiol and 30 μ g/ml *P. mirifica* for 2 and 4 days
 (B) Level of synaptophysin immunoreactivity per 50 μ m neuronal process after treatment of 0.1 μ M 17 β -estradiol and 30 μ g/ml *P. mirifica* for 2 and 4 days
 (C) Densitometric analysis of synaptophysin Western blot of relative intensity for individual experiments was calculated compared to control and averaged
 Immunochemical data represent mean \pm S.E.M. of 5 independent experiments
 *** p < 0.001 vs. control, ** p < 0.01 vs. control, * p < 0.05 vs. control

slip and 3 independent experiments were performed. Western blot was scanned using Scion imaging software and the relative intensity (compared to control) of each independent experiment was evaluated. Data are presented as the mean \pm S.E.M of 5 independent experiments (Western blotting was performed twice in each experiment).

Statistical analysis

All parameters were analyzed using mean \pm standard error of mean (S.E.M.) and student t-test analysis and P value < 0.05 was considered significant.

Results

Effect of *P. mirifica* extract on synaptophysin expression, synaptophysin immunoreactivity in untreated neurons was prominent at puncta whereas they were located mostly along the process in *P. mirifica* and 17 β -estradiol treated groups (Fig. 1A). Administration of *P. mirifica* (30 $\mu\text{g/ml}$) for 4 days significantly increased the level of synaptophysin immunoreactivity along the process in 6-day cultured rat hippocampal neurons ($p < 0.01$ vs control) (Fig. 1B). A similar effect was evident in 17 β -estradiol (0.1 μM) treated group ($p < 0.001$). The quantification of synaptophysin immunoreactivity may not represent the actual protein expression of synaptophysin.

Immunoreactive synaptophysin protein levels quantified by Western blotting indicated that *P. mirifica* (30 $\mu\text{g/ml}$) induced a significant increase in synaptophysin levels after 4 days ($p < 0.05$) but not 2 days of treatment (Fig. 1C). However treatment with 17 β -estradiol (0.1 μM) for 2 and 4 days failed to demonstrate an increase in synaptophysin expression by Western blotting analysis.

The dose-dependency effect of *P. mirifica* on synaptophysin level was tested at 4 doses (3, 15, 30, and 60 $\mu\text{g/ml}$) and a significant stimulatory effect of *P. mirifica* on synaptophysin expression was observed at doses of 30 and 60 $\mu\text{g/ml}$ ($p < 0.05$ and $p < 0.01$ respectively) (Fig. 2). Time-dependent effect of *P. mirifica* (30 $\mu\text{g/ml}$) was evaluated only on the 2nd and 4th day of treatment and synaptophysin was significantly increased on the 4th day after administration ($p < 0.05$) (Fig. 1C, 2, 3).

P. mirifica-induced synaptophysin activation through estrogen receptor

The simultaneous application of the highly specific nuclear estrogen receptor antagonist ICI 182 780 completely prevented the induction effect of *P.*




Fig. 2 Dose-dependent effect of *P. mirifica* on synaptophysin protein level in 6-day cultured rat hippocampal neurons

Densitometric analysis of synaptophysin Western blots were performed after treatment with 3, 15, 30 and 60 $\mu\text{g/ml}$ of *P. mirifica* for 4 days

Data represent the mean \pm S.E.M. of 5 independent experiments

** $p < 0.01$ vs. control, * $p < 0.05$ vs. control




Fig. 3 Effect of ICI 182 780 on the *P. mirifica* induced synaptophysin expression in 6-day cultured rat hippocampal neurons

Densitometric analysis of synaptophysin Western blots were performed after treatment with *P. mirifica* alone (30 $\mu\text{g/ml}$) (*P. mirifica*), ICI 182 780 alone (1 μM) (ICI) and *P. mirifica* plus ICI 182 780 (*P. mirifica* + ICI) for 4 days

Data represent the mean \pm S.E.M. of 5 independent experiments

* $p < 0.05$ vs. control

mirifica (30 $\mu\text{g/ml}$) treatment for 4 days (Fig. 3). ICI 182 780 alone had no influence on synaptophysin level.

Discussion

This is the first report that *P. mirifica* has a significant effect on synaptophysin expression. The fact that *P. mirifica* induced a change in synaptophysin

expression could reflect its effect on synaptogenesis or synapse formation. Synaptophysin was reported to be one of the earliest synaptic proteins to accumulate at developing synapses in culture and its expression is high during synaptogenesis⁽²⁾. The expression of synaptophysin is compatible with its role in regulating synapse formation in cultured hippocampal neurons⁽¹⁹⁾. Also, loss of synaptic proteins, such as synaptophysin and syntaxin, in the frontal neocortex provides the best neurobiological correlates of cognitive impairment⁽²⁰⁾. The effect of *P. mirifica* on synaptophysin expression could be interpreted that it has an indirect effect on the increase in synaptic density associated with neuronal plasticity.

Earlier studies have shown that the administration of estrogen to ovariectomized rats augments protein synthesizing machinery in neurons of the hypothalamic ventromedial nucleus⁽²¹⁾. Estradiol induces rise in spine density and increases dendritic branching in cultured rat hippocampal neurons⁽²²⁾. Phytoestrogen also induces an increase in sexually dimorphic nuclei of preoptic area in male rats⁽²³⁾ and increases aggressive behavior including altered hormonal changes in the brains of male hamsters⁽²⁴⁾. Genistein, a component of soy products, modulates morphological change of gonadotropin-releasing hormone neurons in ovariectomized ewes⁽²⁵⁾, increases the rate of neurite elongation in PC12 cells and produces a concentration-dependent effect on the growth of MCF-7 cells via estrogen receptor⁽²⁶⁾. At a concentration of 50 nM genistein significantly reduces neuronal apoptosis in an estrogen receptor-dependent manner but at a concentration greater than 50 μ M genistein is toxic in primary neurons⁽²⁷⁾. The authors found no significant amount of neuronal death induced by *P. mirifica* (up to 60 μ g/ml). Possible explanations are that the concentration of genistein in *P. mirifica* is lower than the apoptotic dose used in the other studies and *P. mirifica* has no apoptotic effect on hippocampal neurons.

Time course effects of estradiol (0.1 μ g/ml) on spine density production after 4 days of treatment involving alteration of presynaptic and post-synaptic markers were reported⁽²²⁾. This is consistent with our results that the level of synaptophysin after 4 days of treatment with 30 μ g/ml of *P. mirifica* was significantly higher than the level after 2 days of treatment. The results are also in agreement with a recent study demonstrating the ability of estrogen to increase synaptophysin in the CA1 region of the hippocampus in female rhesus macaques (*Macaca mulatta*)⁽⁸⁾.

The nuclear estrogen receptor antagonist,

ICI 182 780, inhibited *P. mirifica*-induced increase in synaptophysin expression in rat hippocampal neurons. ICI 182 780 binds nuclear ER, alters receptor conformation and facilitates binding of co-regulatory proteins that could activate or repress transcriptional activity of estrogen target genes⁽²⁸⁾. It has been reported that phytoestrogen exhibits greater affinity to ER β relative to ER α ⁽²⁹⁾. The present studies are consistent with evidence showing that estrogen-induced increase in dendritic spine density in cultured rat hippocampal neuron is inhibited by ICI 182 780⁽²⁸⁾.

In summary, phytoestrogen-containing *P. mirifica* induced synaptophysin expression in dose- and time-dependent manner in cultured rat hippocampal neurons. Antagonism by ICI 182 780 suggests that this phenomenon may be mediated through estrogen receptor.

Acknowledgements

This work was supported by the Higher Education Development Project of the Commission on Higher Education, Ministry of Education, Thailand.

References

1. Rune GM, Wehrenberg U, Prange-Kiel J, Zhou L, Adelman G, Frotscher M. Estrogen up-regulates estrogen receptor alpha and synaptophysin in slice cultures of rat hippocampus. *Neuroscience* 2002; 113: 167-75.
2. Fletcher TL, Cameron P, De Camilli P, Banker G. The distribution of synapsin I and synaptophysin in hippocampal neurons developing in culture. *J Neurosci* 1991; 11: 1617-26.
3. Papa M, Bundman MC, Greenberger V, Segal M. Morphological analysis of dendritic spine development in primary cultures of hippocampal neurons. *J Neurosci* 1995; 15: 1-11.
4. McEwen BS. Invited review: Estrogens effects on the brain: multiple sites and molecular mechanisms. *J Appl Physiol* 2001; 91: 2785-801.
5. Edelmann L, Hanson PI, Chapman ER, Jahn R. Synaptobrevin binding to synaptophysin: a potential mechanism for controlling the exocytotic fusion machine. *EMBO J* 1995; 14: 224-31.
6. Calakos N, Scheller RH. Vesicle-associated membrane protein and synaptophysin are associated on the synaptic vesicle. *J Biol Chem* 1994; 269: 24534-7.
7. Eastwood SL, Burnet PW, McDonald B, Clinton J, Harrison PJ. Synaptophysin gene expression in human brain: a quantitative in situ hybridization

- and immunocytochemical study. *Neuroscience* 1994; 59: 881-92.
8. Choi JM, Romeo RD, Brake WG, Bethea CL, Rosenwaks Z, McEwen BS. Estradiol increases pre- and post-synaptic proteins in the CA1 region of the hippocampus in female rhesus macaques (*Macaca mulatta*). *Endocrinology* 2003; 144: 4734-8.
 9. Frick KM, Fernandez SM, Bulinski SC. Estrogen replacement improves spatial reference memory and increases hippocampal synaptophysin in aged female mice. *Neuroscience* 2002; 115: 547-58.
 10. Whitten PL, Lewis C, Naftolin F. A phytoestrogen diet induces the premature anovulatory syndrome in lactationally exposed female rats. *Biol Reprod* 1993; 49: 1117-21.
 11. Burton JL, Wells M. The effect of phytoestrogens on the female genital tract. *J Clin Pathol* 2002; 55: 401-7.
 12. Whitehead SA, Cross JE, Burden C, Lacey M. Acute and chronic effects of genistein, tyrphostin and lavendustin A on steroid synthesis in luteinized human granulosa cells. *Hum Reprod* 2002; 17: 589-94.
 13. Suntara A. The remedy pamphlet of Kwao Krua tuber of Luang Anusarnsuntarakromkarnphist. Chiang Mai: Upatipongsa Press; 1931.
 14. Pope GS, Grundy HM, Jone HEH, Tait SAS. The estrogenic substance (miroestrol) from the tuberous roots of *P. mirifica*. *J Endocrinol* 1958; 17: 15-6.
 15. Chansakaow S, Ishikawa T, Sekine K, Okada M, Higuchi Y, Kudo M, et al. Isoflavonoids from *Pueraria mirifica* and their estrogenic activity. *Planta Med* 2000; 66: 572-5.
 16. Malaivijitnond S, Kiatthaipipat P, Cherdshewasart W, Watanabe G, Taya K. Different effects of *Pueraria mirifica*, a herb containing phytoestrogens, on LH and FSH secretion in gonadectomized female and male rats. *J Pharmacol Sci* 2004; 96: 428-35.
 17. Trisomboon H, Malaivijitnond S, Watanabe G, Taya K. Estrogenic effects of *Pueraria mirifica* on the menstrual cycle and hormone-related ovarian functions in cyclic female cynomolgus monkeys. *J Pharmacol Sci* 2004; 94: 51-9.
 18. Bading H, Greenberg ME. Stimulation of protein tyrosine phosphorylation by NMDA receptor activation. *Science* 1991; 253: 912-4.
 19. Tarsa L, Goda Y. Synaptophysin regulates activity-dependent synapse formation in cultured hippocampal neurons. *Proc Natl Acad Sci U S A* 2002; 99: 1012-6.
 20. DeKosky ST, Scheff SW. Synapse loss in frontal cortex biopsies in Alzheimer's disease: correlation with cognitive severity. *Ann Neurol* 1990; 27: 457-64.
 21. Jones KJ, Pfaff DW, McEwen BS. Early estrogen-induced nuclear changes in rat hypothalamic ventromedial neurons: an ultrastructural and morphometric analysis. *J Comp Neurol* 1985; 239: 255-66.
 22. Murphy DD, Segal M. Regulation of dendritic spine density in cultured rat hippocampal neurons by steroid hormones. *J Neurosci* 1996; 16: 4059-68.
 23. Lephart ED, Rhees RW, Setchell KDR, Bu LH, Lund TD. Estrogens and phytoestrogens: brain plasticity of sexually dimorphic brain volumes. *J Steroid Biochem Mol Biol* 2003; 85: 299-309.
 24. Moore TO, Karom M, O'Farrell L. The neuro-behavioral effects of phytoestrogens in male Syrian hamsters. *Brain Res* 2004; 1016: 102-10.
 25. Wojcik-Gladysz A, Romanowicz K, Misztal T, Polkowska J, Barcikowski B. Effects of intracerebroventricular infusion of genistein on the secretory activity of the GnRH/LH axis in ovariectomized ewes. *Anim Reprod Sci* 2005; 86: 221-35.
 26. Wang TT, Sathyamoorthy N, Phang JM. Molecular effects of genistein on estrogen receptor mediated pathways. *Carcinogenesis* 1996; 17: 271-5.
 27. Linford NJ, Dorsa DM. 17beta-Estradiol and the phytoestrogen genistein attenuate neuronal apoptosis induced by the endoplasmic reticulum calcium-ATPase inhibitor thapsigargin. *Steroids* 2002; 67: 1029-40.
 28. Osborne CK, Zhao H, Fuqua SA. Selective estrogen receptor modulators: structure, function, and clinical use. *J Clin Oncol* 2000; 18: 3172-86.
 29. Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH, van der Saag PT, et al. Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology* 1998; 139: 4252-63.